



## If you're using an insulin pump:

- With automated insulin delivery
  - Continue pump and sensor
  - Continue auto mode
  - Do not bolus during prep and prior to the procedure
  - For blood sugars less than 90
    - Exit automated insulin delivery
    - Set a temporary basal rate of 80% until after the procedure is complete
- Without automated insulin delivery
  - Continue pump (and sensor, as indicated)
  - Do not bolus during prep and prior to the procedure
  - For blood sugars less than 90
    - Set a temporary basal rate of 80% until after the procedure is complete
- If your blood sugars remain less than 90 during prep or the morning of the procedure, remove your insulin pump give it to a family member for safe keeping, then reapply when the procedure is complete

## If you're using insulin injections:

- Reduce your long acting insulin by half the night before the procedure
  - Resume your usual dose the next night if eating normally
- Do not use your rapid acting insulin the morning of the procedure
  - Resume your usual dose at the next meal if eating normally

## If you're taking oral medication:

- Hold your medications the morning of the procedure
  - Take your evening medications as scheduled if eating normally
- Resume medications the next morning if eating normally

## Continue your weekly injections as scheduled.

These are general recommendations and are not substitutions for individualized medical advice. Talk to your healthcare provider if you have specific questions or concerns.

**If you have questions or concerns, call the office: 757-659-9903.  
In the event of an emergency, call 911.**